

Warning

Keep out of reach of children

FOR ANIMAL TREATMENT ONLY

VALBAZEN®

Mineralised Cattle

A BENDZIMIDAZOLE (WHITE) DRENCH FOR CATTLE AND RED DEER

Active ingredient: 150g/L Albendazole,

Also contains: copper 68mg and selenium 15mg per 10mL dose

5 Litres

Read entire label before use

HANDLING PRECAUTIONS Warning may be harmful if swallowed, inhaled or absorbed through the skin. May cause mild skin irritation and/or an allergic skin reaction. Suspected of causing genetic defects and may cause organ damage from repeated oral exposure. Danger may damage fertility or the unborn child. Avoid skin contact or inhalation. Wear protective clothing and gloves. Do not breathe vapour.

FIRST AID If **swallowed** call the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. Do NOT induce vomiting. **If on skin** wash with plenty of soap and water. **If inhaled** remove to fresh air. If exposed, feel unwell or a skin irritation/rash occurs, get medical advice.

ENVIRONMENTAL PROTECTION Warning harmful to aquatic organisms with long lasting effects. Harmful to the soil environment. Avoid release to the environment and avoid contamination of any water supply with product or empty container.

DISPOSAL Dispose of any unused contents in an approved landfill. Triple rinse empty container. Recycle or crush/puncture empty container and bury in an approved landfill.

EMERGENCY RESPONSE For specialist advice in an emergency call, 0800 CHEMCALL (0800 243 622)

See Safety Data Sheet for further information. www.spah.co.nz.

DOSE RATE TABLE

Cattle: Roundworms, <i>Ostertagia</i> type II, lungworm and tapeworm			Cattle: Roundworms, <i>Ostertagia</i> type II, lungworm, tapeworm and liver fluke. Red Deer: Roundworms, lungworm.		
LIVEWEIGHT	DOSE	DOSES/PACK	LIVEWEIGHT	DOSE	DOSES/PACK
50kg	2.5mL	2000	60kg	4.0mL	1250
100kg	5.0mL	1000	120kg	8.0mL	625
150kg	7.5mL	666	180kg	12.0mL	416
200kg	10.0mL	500	240kg	16.0mL	312
250kg	12.5mL	400	300kg	20.0mL	250
300kg	15.0mL	333	360kg	24.0mL	208
350kg	17.5mL	285	420kg	28.0mL	177
400kg	20.0mL	250	480kg	32.0mL	156
450kg	22.5mL	222	540kg	36.0mL	138
500kg	25.0mL	200	600kg	40.0mL	125
550kg	27.5mL	181	660kg	44.0mL	113
600kg	30.0mL	166	Over 660kg	1mL/15kg	-
650kg	32.5mL	153			
Over 650kg	1mL/20kg	-			

DIRECTIONS FOR USE

- **SHAKE WELL BEFORE USE.**

- A representative sample of animals should be weighed before treatment.
- Dose rate to be based on the heaviest animal in each group. Do not underdose.
- Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.
- Regularly check the accuracy of drench gun.
- Clean drenching equipment after use.
- Correct drenching technique should be used.

PRECAUTIONS

- Do not use at the same time as any other selenised fertiliser, prill or product and do not exceed the stated dose or dose more frequently than at 3 weekly intervals without consulting a veterinarian.

RESISTANCE

- Resistance can develop to any anthelmintic.
- To delay the onset of resistance:
 - Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your area to reduce development of resistance.
 - It is advisable that a resistance test be conducted regularly when using any parasite treatment.
 - Check effectiveness of anthelmintics at least annually.

EFFICACY

For the treatment and control of susceptible mature and immature roundworms, lungworm in cattle and red deer and mature liver fluke and tapeworm in cattle. Valbazen Mineralised Cattle is effective against inhibited *Ostertagia* Type II and also controls roundworm and fluke eggs in cattle.

At the recommended dosage level, this product is effective for the treatment and control of the following albendazole sensitive parasites:

<u>Common Name</u>	<u>Genus</u>	<u>Species</u>
ROUNDWORMS		
<u>ABOMASUM</u>		
Barber's Pole Worm	<i>Haemonchus</i>	<i>contortus</i>
Small Brown Stomach Worm	<i>Ostertagia</i>	<i>ostertagi</i> <i>leptospicularis</i>
Stomach Hair Worm	<i>Trichostrongylus</i>	<i>axeii</i>
<u>SMALL INTESTINE</u>		
Black Scour Worm	<i>Trichostrongylus</i>	<i>colubriformis</i> <i>vitrinus</i> <i>longispicularis</i>
Small Intestinal Worm	<i>Cooperia</i>	<i>oncophora</i> <i>punctata</i>
Thin-Necked Intestinal Worm	<i>Nematodirus</i>	<i>filicollis</i> <i>spathiger</i> <i>helvetianus</i>
Hookworm	<i>Bunostomum</i>	<i>phlebotomum</i>
<u>LARGE INTESTINE</u>		
Nodule Worm	<i>Oesophagostomum</i>	<i>radiatum</i>
LUNGWORM	<i>Dictyocaulus</i>	<i>viviparus</i>
TAPEWORM	<i>Moniezia</i>	<i>expansa</i>
LIVER FLUKE	<i>Fasciola</i>	<i>hepatica</i>

OVICIDAL

Reduces the output of viable roundworm and fluke eggs.

WITHHOLDING PERIODS

Meat: Cattle and Deer: 7 days – Animals producing meat or offal for human consumption must not be sold for slaughter either during treatment or within seven days of the last treatment.

Milk: 48 hours – Milk intended for sale for human consumption must be discarded during treatment and for not less than 48 hours (i.e. 4 milkings) following the last treatment.

STORAGE

Store at room temperature (ie. 5°C to 25°C).

Registered pursuant to the ACVM Act 1997, No. A6940. See www.nzfsa.govt.nz/acvm/ for registration conditions.

Approved pursuant to the HSNO Act 1996, HSR001941. See www.ermanz.govt.nz for controls

®Pfizer trademark

© Schering-Plough Animal Health Limited, 2007

Registered to:

Schering-Plough Animal Health Limited

33 Whakatiki Street

Upper Hutt, New Zealand

Phone: 0800 800 543

www.spah.co.nz